

### **■ RESEARCH ARTICLE**

# Real world evidence retrospective study to evaluate the safety and efficacy of an amniotic membrane allograft for venous leg ulcers in a post-acute care setting

Ayesha Qadeer,¹ MPH | Emaad Basith,¹ MD | Martha R Kelso,² RN, CHWS, DAPWCA, HBOT | Saad Mohsin,¹ MD | Rashad Sayeed,³ DPM

<sup>1</sup>United WoundCare Institute; <sup>2</sup>Chief Executive Officer for Wound Care Plus, LLC, Kansas City, MO; Head of Research for United Health Partners, LLC, Naperville, IL, US

Correspondence: Ayesha Qadeer (aqadeer@unitedwounds.org)

Received: 23 October 2025 | Accepted: 31 October 2025

Funding: Sponsored by OneBiotech, LLC.

Keywords: Chronic wound | wound healing | acellular and Matrix-like Products (CAMPs) | venous leg ulcer | amniotic

membrane allograft

# **Abstract**

**Aims:** To evaluate the reduction of wound size in venous leg ulcers (VLUs) after amniotic membrane allograft (AmnioBurgeon (OneBiotech, USA)) treatment.

**Methods:** Single-center retrospective database study of patients treated with an amniotic membrane allograft for VLUs. Wound area was measured prior to each allograft application, and the change in wound area after each application was evaluated over the progression of therapy.

**Results:** In total, 38 patients with 43 VLUs were treated weekly with AmnioBurgeon with an average wound size of 9.4 cm<sup>2</sup>. During the treatment course, 6% of the wounds completely healed with a mean time to healing of 38.8 days. 46% of the wounds partially healed with at least 50% or greater area reduction within 17.6 days on average. By the 4th week, 42% of wounds achieved 50% or more reduction in their size. Our treatment group included VLUs ranging between 0.25 cm<sup>2</sup> and 25 cm<sup>2</sup>, achieving significant wound area reduction, indicating wound healing trajectory.

Conclusion: Amniotic membrane allograft is an effective treatment for VLUs.

# Introduction

Wound healing is a complex, multi-phase biological process that proceeds in a defined sequence as the body responds to injury and the surrounding environment.<sup>1</sup> When this process is disrupted for more than four weeks, the wound is classified as chronic and non-healing. Venous leg ulcers (VLUs), which result from chronic venous insufficiency (CVI), are among the most difficult chronic wounds to treat.<sup>2</sup> VLUs represent the most common chronic ulcer type, with a prevalence of 1.5–3 cases per 1,000 individuals and accounting for 50–70% of lower-leg ulcers.<sup>3,4</sup> These wounds contribute to significant morbidity and healthcare expenditure, with an estimated annual cost approaching \$15 billion.<sup>3,4-7</sup> Epidemiological data also indicate higher mortality and poorer outcomes among patients with VLUs.<sup>8</sup>

This is an open access article under the terms of the Creative Commons BY-NC-ND license, which enables reusers to copy and distribute the material in any medium or format in unadapted form only, for noncommercial purposes only, and only so long as attribution is given to the creator.

Several risk factors for VLUs have been identified, many of which are non-modifiable. Advanced age is a major determinant, with prevalence rising to 1.5% among individuals older than 65 years.<sup>6,9</sup> Other contributing factors include prior thromboembolic events, musculoskeletal disorders, obesity, female sex, and genetic predisposition.<sup>3,4,10</sup> The etiology of VLUs is multifactorial involving structural and functional abnormalities of the venous system and failure of the calf-muscle pump, leading to venous hypertension and ulcer formation.<sup>11</sup> Mechanistic contributors include immune dysregulation, alterations in extracellular-matrix remodeling, cytokine imbalance, and aberrant connexin-43.<sup>12</sup> A deeper understanding of these biological pathways may facilitate the development of targeted therapies to improve venous-ulcer healing.<sup>13-16</sup>

Recurrence of VLUs can be caused by various factors such as advanced age, larger wound area, obesity, co-existing venous disease and nutritional deficiencies. <sup>16,17</sup> For each 1cm² in ulcer size, the risk of treatment failure increases by 11%. <sup>17</sup> The standard of care (SOC) for VLUs is compression therapy, which enhances venous return, reduces edema, and optimizes tissue perfusion to support repair. <sup>11</sup> However, compliance remains a challenge due to discomfort, pain, or poor patient understanding of its benefits, underscoring the need for alternative or adjunctive treatments. <sup>2,18-20</sup>

Cellular and tissue-based products (CAMPs) and skin grafting have been increasingly incorporated into VLU management.<sup>21-23</sup> Randomized trials demonstrate that CAMPs significantly enhance wound closure compared with compression alone (SOC), with dehydrated amnion-chorion membranes showing improved healing within defined treatment windows, with dehydrated amnion-chorion membranes showing improved healing within defined treatment windows.<sup>24</sup> However, these studies supported the beneficial role of CAMPs in VLU healing.<sup>21-23,25-28</sup> Apligraf (Organogenesis, USA), an FDA-approved bilayered living skin construct, achieved 47% healing at six months versus 19% with compression alone.<sup>26</sup> CAMPs provide both cellular and matrix support while delivering cytokines and growth factors that promote tissue regeneration.<sup>23,29,30</sup>

Placenta-derived tissues—including amniotic membrane, amniotic fluid, umbilical cord, or composite placental tissue—constitute a category of CAMPs with potent regenerative properties. <sup>23,27-29</sup> These materials promote angiogenesis, modulate inflammation, and limit fibrotic response, thereby accelerating repair. <sup>29</sup> Clinical studies have reported 70% healing at 12 weeks with amniotic-based allografts compared with 30% using SOC alone. <sup>25</sup> Although prior trials have been limited in scale, meta-analyses consistently support improved outcomes among patients treated with amniotic-membrane allografts. <sup>22,27,28</sup> Larger randomized studies remain warranted to further validate these findings.

AmnioBurgeon (OneBiotech, USA) is a resorbable, chorion-free human amnion allograft derived from donated human birth tissue, which has been approved for the management of VLUs. Rich in collagen, growth factors, and extracellular-matrix proteins, AmnioBurgeon supports the biological processes essential to wound repair and has demonstrated favorable outcomes in patients with diabetic foot ulcers (DFUs).

In this retrospective study, we have analyzed the effect of AmnioBurgeon treatment on VLUs.

# Patients and methods

### Data source

The medical charts reviewed in this study spanned the period from March 2025 to September 2025. All treatment records for patients meeting eligibility criteria who received applications of AmnioBurgeon were extracted from the database and de-identified. Atlas 360 electronic medical records were reviewed for the patient's demographic information, clinical history, and laboratory results. Patients' characteristics such as age, gender, weight, comorbidities, and laboratory values such as Ankle-Brachial Index (ABI) were taken. Additional information includes: number of AmnioBurgeon applications, treatment duration, and wound-related examination. Wound area (in cm²) was assessed prior to each treatment wound duration, location, complexity, and signs and symptoms of infection were also documented.

### Ethical approval

For this retrospective study, the Institutional Review Board (IRB), UNIVO, has determined that the study is exempt from IRB oversight requirements under 45 CFR 46.104(d) and under category 4 (#STU25090176). The IRB has approved the request for a waiver of consent and a waiver of authorization for this study.

# Study subjects

The study included medical charts of patients who were treated at United WoundCare Institute in Naperville, IL. Those patients were selected based on eligibility criteria, which are summarized in *Table 1*.

### Outcome and follow-up

The primary goal of this study was to evaluate the change in wound area after AmnioBurgeon treatment for VLUs.

This is an open access article under the terms of the Creative Commons BY-NC-ND license, which enables reusers to copy and distribute the material in any medium or format in unadapted form only, for noncommercial purposes only, and only so long as attribution is given to the creator.

© 2025 The Author(s). International Journal of Tissue Repair

2

### TABLE 1 | Eligibility criteria

Inclusion criteria	Exclusion criteria
<ul> <li>Age 18 years or older</li> <li>Diagnosed with venous ulcer</li> <li>Wound &gt; 4 weeks without significant reduction (&lt;50% healing with 4 weeks of SOC) in wound size</li> </ul>	<ul> <li>Treated with immunosuppressants, corticosteroids, chemotherapy in last 30 days</li> <li>Treated with other cellular, acellular, and matrix-like products (CAMPs) in last 30 days</li> <li>Active infection or cellulitis</li> <li>Active course of antibiotics</li> <li>Lack of baseline or follow-up wound measurements</li> <li>Wounds with a surface area less than 0.25 cm²</li> <li>Wounds with surface area more than 25 cm²</li> <li>Ulcers from other causes such as arterial ulcers, venous ulcers, trauma, surgical, burns</li> </ul>

Complete wound closure was defined as 100% re-epithelialization, as determined by the treating physician. Partial reduction was defined as ≥50% reduction in wound area. No response was defined as <25% reduction in the initial two weeks of treatment or an increase in size. Percentage of wound area reduction, time to reduction, complete wound closure, and number of AmnioBurgeon applications that are required for reduction were also determined. Infections during treatment were assessed by signs and symptoms without the requirement of culture confirmation.

### Statistical analyses

Statistical analyses were conducted using Python (version 3.9.21) and the following packages: Pandas (2.1.4), Numpy (1.26.4), Statsmodels (0.14.0), SciPy (1.11.4), Lifelines (0.28.0), Matplotlib (3.8.2), Seaborn (0.13.0), and the built-in datetime module. Demographics and clinical variables were summarized. Descriptive summaries of continuous variables included the mean, standard deviation, median, and range. Categorical variables were summarized in frequencies and percentages.

A mixed-effects linear regression model accounting for repeated measures was performed to assess significance (p<0.05) in the change of wound area before and after treatment, and between applications. Baseline wound size areas were placed into two groups (<10 cm², and 10 to <24.99 cm²). Wounds were separated based on age into 2 groups: <180 days or ≥180 days. Chi-squared test was performed to compare the proportion of wounds achieving ≥50% area reduction after 28 days of treatment between groups. Kaplan-Meier analysis was performed to compare the time to ≥50% wound area reduction between groups. Recurrent wounds were excluded from analysis to avoid over-representation.

### Results

### Patients' demographics and wound characteristics

In this study, 47 VLUs in 38 patients were treated with AmnioBurgeon after failing to heal with SOC therapy. The median age of the patients was 79.6 (range 48–100), and 47% were female (*Table 2*). At baseline, the median wound area was 8.8 cm² (range 1.2–20.3). The median wound duration prior to CAMP treatment was 180 days, and most wounds have their fat layers exposed (~39%). Wounds were treated with CAMP on average every 7.4±1.1 days, as literature suggests, reflecting both the biologic needs of wound healing and the realities of patient care delivery. The median number of CAMP applications per wound was 6 (range 2-12). The median duration of the CAMP treatment course was 35 days (7-96).

# Wound healing efficacy outcome

AmnioBurgeon treatment resulted in an overall favorable outcome in VLUs that failed SOC treatment (*Table 3*). Of the 47 wounds, 37 (46%) wounds achieved more than ≥50% reduction in wound area. The median time to ≥50% reduction was 14 days (range 7, 49) and required a median of 6 (range 2, 12) CAMP applications. At least 50% wound area reduction after 28 days of treatment is a strong predictor of potential complete wound closure, which was observed in 20 (42%) wounds. This result showed a substantial early response to the treatment. AmnioBurgeon treatment resulted in complete wound closure in 6% of wounds. The median time to complete wound closure was 42 days (range 14-72) and required a median of 4.5 applications. No response (<25% reduction or increased size) was noted in 14 (29%) wounds.

### Baseline wound size

Baseline wound size impacted the percent area reduction with the number of CAMP treatment applications (*Figure 1*). The PAR effect of more CAMP applications for wound sizes <10cm² was more pronounced than for wounds between 10 and 25 cm².

This is an open access article under the terms of the Creative Commons BY-NC-ND license, which enables reusers to copy and distribute the material in any medium or format in unadapted form only, for noncommercial purposes only, and only so long as attribution is given to the creator.

TABLE 2 | Patient demographics and wound characteristics

1	
Number of patients	38
Age, years (Mean±SD, Median, Min, Max)	79.03 ±10.8, 79.6, 48.8, 100.24
Female (N, %)	18 (47%)
Number of wounds	47
Wound locations	
Lower leg	35
Feet	12
Stage/severity	
Fat layer exposed	15
Breakdown of Skin	6
Unspecified severity	25
Muscle involvement without evidence of necrosis	1
Without bone or tendon exposure	0
Without evidence of necrosis	0
Wound size at baseline (cm²) (Mean±SD, Median, Min, Max)	9.4±5.1, 8.8, 1.2, 20.3
Wound Age (Days) Median	180
Time from First Visit to CAMPs Treatment, (Days) (Mean±SD, Median)	16.8±31.5, 126.0

Additionally, baseline wound size was shown to affect the likelihood of achieving  $\geq$ 50% reduction in wound area after treatment (*Figure 2*). Kaplan-Meier analysis with log-rank test was conducted to show wounds <10 cm², 53.8% achieved partial healing at 22 days, while 42.9% of wounds sized between 10-24.99 cm² achieved partial healing in 49 days. Findings showed that smaller wounds trended towards faster healing, but the difference was not statistically significant (p=0.68).

Wound age was found to significantly affect the response to AmnioBurgeon treatment (Figure~3). Wounds aged <180 days (n=22) reached partial reduction by day 14 in 72.7% (n=16) wounds compared to only 28% (n=7) in wounds with age >= 180 days (n=25). Wounds aged <180 days demonstrated a statistically significantly shorter time to achieving partial healing (log-rank p=0.0018).

# Discussion

VLUs are a debilitating healthcare problem with significant morbidity and costs.<sup>5</sup> The leading underlying pathology for VLUs is chronic venous insufficiency, and this forms the basis and emphasizes the necessity of compression therapy in the treatment of VLUs.<sup>33</sup> SOC for VLUs, including previously mentioned compression therapy, dressing, and others. However, the healing rate is still unsatisfactory. Several factors have been identified as contributing to treatment failure. This includes advanced age, which is associated with less mobility; a factor known to aggravate the underlying pathology, while other risk factors include wound age.

The use of CAMPs in the treatment of chronic wounds, including VLUs, is growing. Skin grafting is a primary therapy for wounds larger than 25cm². Several studies have demonstrated higher closure rates with amniotic-based CAMPs. Despite the results being promising, larger-scale studies are needed. In our study, we analyzed retrospective data of patients treated with AmnioBurgeon for their hard-to-heal VLUs that were unsuccessful with SOC alone. Our results were encouraging despite several challenging factors in the baseline patient population, such as advanced age (mean 79.6), patient compliance, and other ongoing comorbidities.

In this retrospective analysis, we studied the effect of AmnioBurgeon on VLU healing. AmnioBurgeon is a resorbable, chorion-free human amnion allograft derived from donated human birth tissue, which has been approved for patients

This is an open access article under the terms of the Creative Commons BY-NC-ND license, which enables reusers to copy and distribute the material in any medium or format in unadapted form only, for noncommercial purposes only, and only so long as attribution is given to the creator.

<sup>© 2025</sup> The Author(s). International Journal of Tissue Repair

### TABLE 3 | Wound healing treatment outcomes

Completely Healed Wound (%)	3 (6%)
Complete healing time (days) (Mean±SD, Median, Min, Max)	38.8±19.6, 42.0, 14.0, 72.0
Number of CAMPs Applications for Completely Healed Wound (Mean±SD, Median, Min, Max)	4.2±1.6, 4.5, 2.0, 6.0
Partial Healed Wound (>=50%)	22 (46%)
Partial Healed time (days) (Mean±SD, Median, Min, Max)	17.6±12.4, 14.0, 7, 49
>=50% reduction by day 28 (%)	20 (42%)
Number of CAMP Applications for Partial Healed Wound (Mean±SD, Median, Min, Max)	6.4±2.9, 6.0, 2, 12
No Response (<25% reduction or increased size) (%)	14 (29%)
Duration of CAMPs Applications (Mean±SD, Median, Min, Max)	39.1±23.5, 35.0, 7, 96
Number of CAMPs Applications (Mean±SD, Median, Min, Max)	6.2±3.0, 6.0, 2, 12
Time until Max PAR (Mean±SD, Median, Min, Max)	39.10, 23.53, 35, 7, 96
Duration between CAMPs Applications (days) (Mean±SD, Median, Min, Max)	7.4±1.1, 7.0, 6, 14
ABI (n, Mean, Median, Min, Max)	47, 0.887234, 0.86, 0.6, 1.32
Infection (n, %)	13 (34%)

# PAR vs Application Number by Wound Size Group




FIGURE 1 | In a mixed-effects linear regression model accounting for repeated measures within wounds, each additional Cellular, Acellular, Matrix-like Product (CAMPs) application was associated with a significant increase in percent area reduction for smaller wounds (PAR) (p=0.068).

This is an open access article under the terms of the Creative Commons BY-NC-ND license, which enables reusers to copy and distribute the material in any medium or format in unadapted form only, for noncommercial purposes only, and only so long as attribution is given to the creator.

<sup>© 2025</sup> The Author(s). International Journal of Tissue Repair

# Partial Healing (>=50%) by Wound Size Group (Kaplan-Meier) Log-rank test: Chi-sq = 0.17, p = 0.6765 0.1 Probability Wound Not Partially Healed 0.8 9.0 Area Group <10cm<sup>2</sup> 10-24.99cm<sup>2</sup> 0.4 0.2 0.0 0 10 20 30 40 60 50 Days to Partial Heal (PAR >= 50%)

FIGURE 2 | Kaplan-Meier analysis across wound size groups to time of partial healing. Description: Time to partial healing (≥50% PAR) by wound size. Survival curves showing time until wounds achieve ≥50% area reduction, comparing wounds that are (<10cm²) vs (10-24.99cm²). Key findings: Median time to partial healing <10cm² (53.8%) wounds in 22 days, while 10-24.99cm² (42.9%) wounds in 49 days. Statistical test: Log-rank test: x2=0.17, p=068 (not significant). Conclusion: Smaller wounds exhibited a trend toward faster healing (22 vs 49 median days), but the difference was NOT statistically significant. The sample size may be too small to detect a true difference, or there is substantial overlap in healing times between the two groups.




FIGURE 3 | Kaplan-Meier analysis with a log-rank test demonstrated a statistically significant difference in time to partial healing between wounds aged <180 days and ≥180 days (log-rank p = 0.0018), with younger wounds achieving partial healing more rapidly.

This is an open access article under the terms of the Creative Commons BY-NC-ND license, which enables reusers to copy and distribute the material in any medium or format in unadapted form only, for noncommercial purposes only, and only so long as attribution is given to the creator.

with VLU. We measured the effect of AmnioBurgeon across 47 wounds that failed SOC treatment. AmnioBurgeon was found to result in complete wound healing in 6% of wounds and wound size reduction by 50% or more at 4 weeks in 42% of patients. This is a strong predictor of complete healing at 12 weeks,<sup>34-36</sup> and it serves as a valid endpoint in chronic wound management. Performing the same tests, we found a significant difference in time to partial healing between wounds aged less than 180 days and those with more than 180 days. This may further suggest that wounds that have been ongoing for more than 6 months are already very resistant and associated with poorer outcomes.<sup>37</sup>

# Study limitations

This study has several limitations. It was a single-center, retrospective analysis with a relatively small sample size, which may limit the generalizability of the findings. The study design inherently lacked randomized controls, making it difficult to account for potential confounding variables or to establish causality. Additionally, variability in patient compliance with compression therapy and follow-up visits may have influenced healing outcomes. The short duration of observation also restricts the assessment of long-term wound closure rates and recurrence. Future larger, multicenter, and prospective studies with extended follow-up and standardized treatment protocols are needed to validate these findings.

### Conclusion

Given the complexity and systemic nature of VLUs, they remain among the most challenging chronic wounds to treat. Despite these challenges, AmnioBurgeon has demonstrated safety and effectiveness as a CAMP therapy, promoting rapid wound size reduction. Future studies evaluating larger patient populations will be necessary.

# Acknowledgments

Support in data collection, cleaning, validation and visualization: Mohsin Ansari; Hafsa Ahmed; Maham Khan; Maham Surtee; Naz Khan.

### Conflicts of interest

The authors have no conflicts of interest to declare.

### Data availability statement

Data is not publicly available.

# Ethics approval and consent to participate

UNIVO IRB approval.

### **Author contributions**

Ayesha Qadeer: Study design, data analysis and authorship; Emaad Basith: Medical expert and authorship; Martha R Kelso: Medical expert and authorship; Saad Mohsin: Medical expert and authorship; Rashad Sayeed: Study design, medical expert and authorship.

# References

- 1. Sen CK. Human Wounds and Its Burden: An Updated Compendium of Estimates. Adv Wound Care (New Rochelle). 2019;8(2):39-48. https://doi.org/10.1089/wound.2019.0946
- Mostow EN, Haraway GD, Dalsing M, Hodde JP, King D; OASIS Venus Ulcer Study Group. Effectiveness of an extracellular matrix graft (OASIS Wound Matrix) in the treatment of chronic leg ulcers: a randomized clinical trial. J Vasc Surg. 2005;41(5):837-843. https://doi.org/10.1016/j.jvs.2005.01.042
- 3. Bonkemeyer Millan S, Gan R, Townsend PE. Venous Ulcers: Diagnosis and Treatment. Am Fam Physician. 2019;100(5):298-305.
- Gillespie DL; Writing Group III of the Pacific Vascular Symposium 6, Kistner B, et al. Venous ulcer diagnosis, treatment, and prevention of recurrences. J Vasc Surg. 2010;52(5 Suppl):8S-14S. https://doi.org/10.1016/j. jvs.2010.05.068
- Olin JW, Beusterien KM, Childs MB, Seavey C, McHugh L, Griffiths RI. Medical costs of treating venous stasis ulcers: evidence from a retrospective cohort study. Vasc Med. 1999;4(1):1-7. https://doi. org/10.1177/1358836X9900400101
- 6. Falanga V, Isseroff RR, Soulika AM, et al. Chronic wounds. Nat Rev Dis Primers. 2022;8(1):50. Published 2022 Jul 21. https://doi.org/10.1038/s41572-022-00377-3
- 7. Xie T, Ye J, Rerkasem K, Mani R. The venous ulcer continues to be a clinical challenge: an update. Burns Trauma. 2018;6:18. https://doi.org/10.1186/s41038-018-0119-y
- 8. Salenius JE, Suntila M, Ahti T, et al. Long-term Mortality among Patients with Chronic Ulcers. Acta Derm Venereol. 2021;101(5):adv00455. https://doi.org/10.2340/00015555-3803
- 9. Margolis DJ, Bilker W, Santanna J, Baumgarten M. Venous leg ulcer: incidence and prevalence in the elderly. J Am

This is an open access article under the terms of the Creative Commons BY-NC-ND license, which enables reusers to copy and distribute the material in any medium or format in unadapted form only, for noncommercial purposes only, and only so long as attribution is given to the creator.

- Acad Dermatol. 2002;46(3):381-386. https://doi.org/10.1067/mjd.2002.121739
- 10. Grzela T, Bialoszewska A. Genetic risk factors of chronic venous leg ulceration: Can molecular screening aid in the prevention of chronic venous insufficiency complications?. Mol Med Rep. 2010;3(2):205-211. https://doi.org/10.3892/mmr 00000241
- 11. Patton D, Avsar P, Sayeh A, et al. A systematic review of the impact of compression therapy on quality of life and pain among people with a venous leg ulcer. Int Wound J. 2024;21(3):e14816. https://doi.org/10.1111/iwj.14816
- 12. Kirsner RS, Baquerizo Nole KL, Fox JD, Liu SN. Healing refractory venous ulcers: new treatments offer hope. J Invest Dermatol. 2015;135(1):19-23. https://doi.org/10.1038/jid.2014.444
- 13. Chi YW, Raffetto JD. Venous leg ulceration pathophysiology and evidence based treatment. Vasc Med. 2015;20(2):168-181. https://doi.org/10.1177/1358863X14568677
- 14. Crawford JM, Lal BK, Durán WN, Pappas PJ. Pathophysiology of venous ulceration. J Vasc Surg Venous Lymphat Disord. 2017;5(4):596-605. https://doi.org/10.1016/j.jvsv.2017.03.015
- 15. Malanin K. About the pathophysiology of venous leg ulceration. J Am Acad Dermatol. 2002;47(1):157-158. https://doi.org/10.1067/mjd.2002.120577
- Raffetto JD, Ligi D, Maniscalco R, Khalil RA, Mannello F. Why Venous Leg Ulcers Have Difficulty Healing: Overview on Pathophysiology, Clinical Consequences, and Treatment. J Clin Med. 2020;10(1):29. https://doi.org/10.3390/ icm10010029
- 17. Margolis DJ, Berlin JA, Strom BL. Risk factors associated with the failure of a venous leg ulcer to heal. Arch Dermatol. 1999;135(8):920-926. https://doi.org/10.1001/archderm.135.8.920
- 18. Stevenson EM, Coda A, Bourke MDJ. Investigating low rates of compliance to graduated compression therapy for chronic venous insufficiency: A systematic review. Int Wound J. 2024;21(4):e14833. https://doi.org/10.1111/iwj.14833
- 19. Weller CD, Richards C, Turnour L, Team V. Patient Explanation of Adherence and Non-Adherence to Venous Leg Ulcer Treatment: A Qualitative Study. Front Pharmacol. 2021;12:663570. https://doi.org/10.3389/fphar.2021.663570
- Bar L, Marks D, Brandis S. Developing a Suite of Resources to Improve Patient Adherence to Compression Stockings: Application of Behavior Change Theory. Patient Prefer Adherence. 2023;17:51-66. https://doi. org/10.2147/PPA.S390123
- 21. Jones JE, Nelson EA, Al-Hity A. Skin grafting for venous leg ulcers. Cochrane Database Syst Rev. 2013;2013(1):CD001737. https://doi.org/10.1002/14651858.CD001737.pub4
- 22. Pan J, Hu X, Yin H, Zhang C, Yan Z. Effectiveness of different types of skin grafting for treating venous leg ulcers: A protocol for systematic review and network meta-analysis. Medicine (Baltimore). 2021;100(15):e25597. https://doi.org/10.1097/MD.00000000000025597
- 23. Deng X, Gould M, Ali MA. A review of current advancements for wound healing: Biomaterial applications and medical devices. J Biomed Mater Res B Appl Biomater. 2022;110(11):2542-2573. https://doi.org/10.1002/jbm.b.35086
- 24. Bianchi C, Cazzell S, Vayser D, et al. A multicentre randomised controlled trial evaluating the efficacy of dehydrated human amnion/chorion membrane (EpiFix®) allograft for the treatment of venous leg ulcers. Int Wound J. 2018;15(1):114-122. https://doi.org/10.1111/iwj.12843
- 25. Serena TE, Orgill DP, Armstrong DG, et al. A Multicenter, Randomized, Controlled, Clinical Trial Evaluating Dehydrated Human Amniotic Membrane in the Treatment of Venous Leg Ulcers. Plast Reconstr Surg. 2022;150(5):1128-1136. https://doi.org/10.1097/PRS.00000000000009650
- 26. Falanga V, Sabolinski M. A bilayered living skin construct (APLIGRAF) accelerates complete closure of hard-to-heal venous ulcers. Wound Repair Regen. 1999;7(4):201-207. https://doi.org/10.1046/j.1524-475x.1999.00201.x
- 27. Alomairi AA, Alhatlani RA, Alharbi SM, et al. Assessing the application and effectiveness of human amniotic membrane in the management of venous and diabetic ulcers: a systematic review and meta-analysis of randomized controlled trials. Cureus. 2024;16(3):e56659. Published 2024 Mar 21. https://doi.org/10.7759/cureus.56659
- 28. Shaydakov ME, Ting W, Sadek M, et al. Review of the current evidence for topical treatment for venous leg ulcers. J Vasc Surg Venous Lymphat Disord. 2022;10(1):241-247.e15. https://doi.org/10.1016/j.jvsv.2021.06.010
- 29. Protzman NM, Mao Y, Long D, et al. Placental-Derived Biomaterials and Their Application to Wound Healing: A Review. Bioengineering (Basel). 2023;10(7):829. https://doi.org/10.3390/bioengineering10070829
- 30. Vecin NM, Kirsner RS. Skin substitutes as treatment for chronic wounds: current and future directions. Front Med (Lausanne). 2023;10:1154567. https://doi.org/10.3389/fmed.2023.1154567
- 31. Zelen CM, Serena TE, Snyder RJ. A prospective, randomised comparative study of weekly versus biweekly application of dehydrated human amnion/chorion membrane allograft in the management of diabetic foot ulcers. Int Wound J. 2014;11(2):122-128. https://doi.org/10.1111/iwj.12242
- 32. Lavery LA, Suludere MA, Raspovic K, Crisologo PA, Johnson MJ, Tarricone AN. Randomized Controlled Trial to Compare AmnioExcel Human Amniotic Allograft in Weekly Versus Biweekly Treatment of Diabetic Foot Ulcers. Int J Low Extrem Wounds. https://doi.org/10.1177/15347346241276697
- 33. Harding K. Challenging passivity in venous leg ulcer care the ABC model of management. Int Wound J. 2016;13(6):1378-1384. https://doi.org/10.1111/iwj.12608
- 34. Serena TE, Yaakov R, DiMarco D, et al. Dehydrated human amnion/chorion membrane treatment of venous leg

- ulcers: correlation between 4-week and 24-week outcomes. J Wound Care. 2015;24(11):530-534. https://doi.org/10.12968/jowc.2015.24.11.530
- 35. Gwilym BL, Mazumdar E, Naik G, Tolley T, Harding K, Bosanquet DC. Initial Reduction in Ulcer Size As a Prognostic Indicator for Complete Wound Healing: A Systematic Review of Diabetic Foot and Venous Leg Ulcers. Adv Wound Care (New Rochelle). 2023;12(6):327-338. https://doi.org/10.1089/wound.2021.0203
- 36. Masiello G, Franzini M, Tirelli U, et al. Successful treatment of severe venous leg ulcers and diabetic foot ulcers using ozone. J Vasc Surg Venous Lymphat Disord. 2025;13(6):102278. https://doi.org/10.1016/j.jvsv.2025.102278
- 37. Fife CE, Horn SD. The Wound Healing Index for Predicting Venous Leg Ulcer Outcome. Adv Wound Care (New Rochelle). 2020;9(2):68-77. https://doi.org/10.1089/wound.2019.1038